

Poster presentation

HAART in HIV+ naive elderly patients: immuno-virological response and clinical outcome

BM Celesia*¹, F Bisicchia¹, R La Rosa², S Mavilla¹, M Gussio¹, MT Mughini¹, F Palermo¹ and R Russo¹

Address: ¹Unit of Infectious diseases University of Catania ARNAS Garibaldi, Catania, Italy and ²Unit of Infectious diseases Ferrarotto Hospital AO Vittorio Emanuele, Catania, Italy

* Corresponding author

from Ninth International Congress on Drug Therapy in HIV Infection
Glasgow, UK. 9–13 November 2008

Published: 10 November 2008

Journal of the International AIDS Society 2008, **11**(Suppl 1):P17 doi:10.1186/1758-2652-11-S1-P17

This abstract is available from: <http://www.jiasociety.org/content/11/S1/P17>

© 2008 Celesia et al; licensee BioMed Central Ltd.

Purpose of the study

Elderly patients (≥ 50 years) are increasing in the HIV population; HAART-related prolonged survival and late diagnosis of new HIV infections are possible reasons of this findings. It is debated whether older patients have a different response to HAART. The aim of this retrospective study was to evaluate efficacy of HAART and clinical outcome in a group of patients ≥ 50 year in comparison to a control group (< 50 years-old).

Methods

All naive patients starting HAART since November 1996 in two different Infectious Diseases units in Catania (Sicily) were included. The following parameters were evaluated: epidemiological (sex, age, risk factors, year of HAART beginning), viro-immunological (CD4 cell count, HIV-RNA viral load), and clinical (CDC, first treatment, number and causes of therapeutic switch, new AIDS diseases and death). The follow-up was performed until the last available visit.

Summary of results

We enrolled 276 HIV-1-infected patients; 215 (78%) male, median age 38 years; 122 (44%) heterosexuals, 103 (38%) homo-bisexuals, 43 (16%) drug addicts; 139 (50.4%) CDC A, 31 (11.2%) CDC B, 106 (38.4%) CDC C; median CD4 cell count was 155 (IQ range 48–301), median HIV-RNA viral load 5.0 log₁₀ (IQ range 4.3–5.4). Fifty-one patients (18.5%) were ≥ 50 years old, 82.4%

male, most of them infected by sexual intercourse ($p = 0.002$). At baseline, elderly patients were more frequently symptomatic ($p = 0.002$) with a trend to lower CD4 cell count (98 vs. 169 cells/ μ l); no differences were seen on HIV-RNA copies/ml. Twelve months after beginning of HAART, median absolute increase (146 vs. 165 cells/ μ l) of the CD4 cell count and percentage of patients with HIV-RNA < 400 copies/ml (87.2% vs. 85.7%) were comparable in elderly and younger patients, respectively 82.1% and 82.8% achieved an immuno-virological response (defined as VL < 400 copies/ml and more than 100 CD4 cells/ μ l increase). At last, 65% and 16% of elderly patients achieved, respectively, more than 200 and more than 500 CD4 cell/ μ l with no significant difference with younger patients. Discontinuation of HAART (any causes) was less frequent in older subjects ($p = 0.04$). After a median follow-up of 68 months, the percentage of deaths was 15.7% and 5.8% ($p = 0.034$) in the elderly and younger group, respectively.

Conclusion

In our experience, elderly and younger naive patients on HAART have similar immunological and virological response while hard clinical end-points tend to be more frequent in older subjects. Prospective studies are necessary to further investigate our findings.